



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

Syllabus

Remark 1.

The law of large numbers

Let be a series of independent two by two random variables X_1, X_2, \dots, X_n , which admit finite mean values and bounded dispersions:

X_1, X_2, \dots, X_n a.i. $M(X_k) < \infty \forall k \in N^*, D^2(X_k) \leq c < \infty, \forall k \in N^*, c > 0$ then:

$$(1) \quad \left[\frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{\text{almost sure}} \frac{1}{n} \sum_{k=1}^n M(X_k) \right], \quad \lim_{n \rightarrow \infty} P \left(\left| \frac{1}{n} \sum_{k=1}^n X_k - \frac{1}{n} \sum_{k=1}^n M(X_k) \right| < \varepsilon \right) = 1, \forall k \in N^* \quad (2)$$

Particular cases:

A. If the random variables have the same mean: $M(X_k) = m, \forall k \in N^*, \frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{\text{almost sure}} m$

B. If the random variables are *Bernoulli type*:

$$X_k = \begin{pmatrix} 1 & 0 \\ p & q \end{pmatrix}, p, q > 0, p + q = 1 \Rightarrow M(X_k) = p, \forall k \in N, \frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{\text{almost sure}} p \quad (3) \Leftrightarrow f_n = \frac{1}{n} \sum_{k=1}^n X_k, f_n \xrightarrow{\text{a.s.}(n \rightarrow \infty)} p$$

Central-limit theorem

Let be $(X_n)_{n \in N^*}$ a series of independent two by two random variables, which admit finite mean values and bounded dispersions:

$$\frac{\sum_{k=1}^n X_k - \sum_{k=1}^n M(X_k)}{\sqrt{\sum_{k=1}^n D^2(X_k)}} \xrightarrow{\text{repartition}(n \rightarrow \infty)} N(0,1).$$




Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SAN JORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

Remark 2.

$X \in N(0,1)$, is the random variable having a normalized normal distribution if the *distribution density* is defined by:

$$\varphi_X(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}, \forall x \in R.$$

Its cumulative distribution function is:

$$F(x) = P(X < x) = \int_0^x f(t) dt = \frac{1}{\sqrt{2\pi}} \int_0^x e^{-t^2/2} dt = \phi(x), \forall x \in R.$$

where $\phi(x)$ is *Laplace function* having the following properties:

$$\phi(0) = 0, \text{ functie tabelata}; \phi(-x) = 1 - \phi(x); \phi(\infty) = \frac{1}{2}, \phi(-\infty) = \frac{1}{2}.$$

Error function is defined as $\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt, \forall x \in R$ and express the probability of a random variable, with normal distribution of mean 0 and variance 1/2 falling in the range $[-x, x]$. $\phi(x) = \frac{1}{2} [1 + \text{erf}(\frac{x}{\sqrt{2}})]$.

Remark 3.

From the analytical form of the normal distribution function results the impossibility of explicitly solving an equation of form $F(x)=y$. This leads to the development of special methods for evaluating the solution $x = F^{-1}(y)$. One of these methods is based on the Proposition:

Proposition 1

If U and V are 2 independent random variables, uniform distributed in the interval $[0,1]$, then the random variables:

$$X = \sqrt{-2 \cdot \ln(U)} \cdot \cos(2\pi \cdot V); Y = \sqrt{-2 \cdot \ln(U)} \cdot \sin(2\pi \cdot V)$$

are independent and normal distributed, with mean $m = 0$ and mean square deviation $\sigma^2=1$.




Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SAN JORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

For Z normal standard distribution, $X = \sigma Z + m$ is the cumulative normal distribution function with mean $m \neq 0$ and $\sigma \neq 1$:


$$P(m - k\sigma < X < m + k\sigma) = P(-k < Z < k) = 2\Phi(k) - 1$$

For $k=3$ we have:

$P[m - 3\sigma < X < m + 3\sigma] = 0.997$, meaning that 99,7% values of the function Z are in the interval.

Examples of distributions functions

1. Normal distribution with parameters m and σ^2 , $X \sim N(m, \sigma^2)$; $X \approx N(m, \sigma^2) \Rightarrow \frac{X - m}{\sigma} \sim N(0, 1)$



2. Uniform distribution $X \sim U[a, b]$

X v.a. with density function: $f_X(x) = \begin{cases} \frac{1}{b-a}, & x \in [a, b] \\ 0 & \text{in rest} \end{cases}$. X has the mean $M(X) = (a + b)/2$ and dispersion $D(X) = (b - a)^2 / 12$.



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

3. Exponential distribution $X \sim \text{Exp}(\lambda, \alpha)$


X r.d. with *density function* funcția: $f_X(x) = \begin{cases} \frac{1}{\lambda} e^{-\frac{x-\alpha}{\lambda}}, & x \geq \alpha, \lambda > 0 \\ 0 & \text{în rest} \end{cases}$.

X has the mean $M(X) = \lambda + \alpha$ and dispersion $D(X) = \lambda^2$. Remark: $\text{Exp}(\lambda, 0) = \text{Exp}(\lambda)$.

Negative exponential distribution


X has *density function*: $f(x) = \lambda e^{-\lambda x}$, $x \in [0, \infty)$, $\lambda > 0$,

or cdf $F(x) = P[X < x] = \begin{cases} 1 - e^{-\lambda x}, & x \geq 0 \\ 0, & \text{otherwise} \end{cases}$.



4. Weibull distribution $X \sim W(\alpha, \beta)$

X r.d. having *density function*: $f_X(x) = \alpha \beta x^{\beta-1} e^{-(\alpha x^\beta)}$
with mean $M(X) = \lambda + \alpha$ and dispersion $D(X) = \lambda^2$.





Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SAN JORGE




ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

5. chi-square distribution $X \sim \chi^2(n)$

X r.d. with *density function*: $f_X(x) = \begin{cases} \frac{x^{n/2-1}}{2^{n/2}\Gamma(\frac{n}{2})} e^{-x/2}, & x \geq 0, \\ 0 & \text{in rest} \end{cases}$, and $\Gamma(p) = \int_0^\infty x^{p-1} e^{-x} dx$.




6. Student distribution $X \sim T(n)$ with n degrees of freedom

X r.d. with *density function*: $f_X(x) = \frac{\Gamma(\frac{n+1}{2})}{\sqrt{\pi \cdot n} \Gamma(\frac{n}{2})} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}$.

Proposition:

For U, V two random variables such that $U \sim N(0,1)$ and $V \sim \chi^2$ then:

$$W = U / \sqrt{V/n} \cdot \sim T(n)$$





Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE




ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

7. Fisher distribution with n , and m degrees of freedom $X \sim F(n, m)$

X r.d. with *density function*

$$f_X(x) = \frac{\Gamma\left(\frac{m+n}{2}\right)}{\Gamma\left(\frac{m}{2}\right)\Gamma\left(\frac{n}{2}\right)} \left(\frac{n}{m}\right)^{n/2} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}$$




8. Beta distribution $X \sim \text{Be}(\alpha, \beta)$, $\alpha > 0, \beta > 0$

X r.d. with *density function*:

$$f_X(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{B(\alpha, \beta)}, x \in (0, 1), \text{ cu } B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}.$$

$$M(X) = \frac{a}{a+b}, \text{ Var}(X) = \frac{ab}{(a+b)^2(a+b+1)}$$



9. Gamma distribution $G(a, b)$, $a > 0, b > 0$.

X r.d. with *density function*: $f_X(x) = \frac{x^{b-1}e^{-ax}a^b}{\Gamma(b)}$, $x > 0$ with $M(X) = b/a$; $\text{Var}(X) = b/a^2$.

For $b=1$ one obtain exponential distribution $\text{Exp}(a)$.



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SAN JORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES


Discrete distributions

Binomial distribution $X \sim Bi(p, n)$.

$$P(X = k) = C_n^k p^k (1 - p)^{n-k}$$

$$M(X) = np; \quad \text{Var}(X) = npq.$$

represents the probability with which an event A is realized k times in a number n (fixed) of samples in which the probability of realization of A in each sample is p.



Negative Binomial distribution

$$X \sim NB(r, p)$$

$$P(X = k) = C_{r+k-1}^k p^r q^k; \quad k=0, 1, 2, \dots$$

$$M(X) = r \frac{q}{p}; \quad \text{Var}(X) = r \frac{q}{p^2}.$$


is the probability of having k failures before the second success in a series of Bernoulli tests (the result of each test consists in the realization of an event A with the probability p or of the event opposite to it with the probability q).

Poisson distribution

$$X \sim Po(\lambda). \quad P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}$$

$$M(X) = \text{Var}(X) = \lambda.$$

express the number of events that result in property damage.





Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

The Moment generating Function (MGF)

Definition :

The moment generating function (mgf) of a random variable X is given by:

$$M_X(t) = E[e^{tX}] \quad , \quad t \in \mathbb{R}.$$

Theorem!

If the random variable X has the (mgf) $M_X(t)$ then:

$$M_X^{(r)}(0) = \left. \frac{d^r}{dt^r} [M_X(t)] \right|_{t=0} = E(X^r).$$

The derivative of order r of the (mgf) evaluated at $t=0$ gives the value of the r^{th} moment.



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SAN JORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

Remark:

For a discrete probability mass function $M_X(t) = \sum_{i=1}^{\infty} e^{tx_i} p_i$

For a continuous probability density function, $M_X(t) = \int_{-\infty}^{\infty} f(x) e^{tx} dx$.

Theorem 2:

Let X be a random variable with $M_X(t)$ the (mgf) and a, b constants. If $Y = a + bX$ then (mgf) of Y is given by

$$M_Y(t) = e^{at} M_X(bt)$$



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

Example 1.

$X \sim \text{Po}(\lambda)$ with probability mass function $p(x) = \frac{e^{-\lambda} \lambda^x}{x!}$, $x=0,1,\dots$

$$M_X(t) = E[e^{tx}] = \sum_{x=0}^{\infty} e^{tx} \frac{e^{-\lambda} \lambda^x}{x!} = e^{-\lambda} \sum_{x=0}^{\infty} \frac{(e^t \lambda)^x}{x!}$$

$$M_X(t) = e^{-\lambda} e^{e^t \lambda} = e^{\lambda(e^t - 1)}$$

$$\text{Thus: } M_X'(t) = \frac{d}{dt} [e^{\lambda(e^t - 1)}] = \lambda e^t e^{\lambda(e^t - 1)}$$

$$M_X''(t) = \frac{d^2}{dt^2} [e^{\lambda(e^t - 1)}] = \lambda e^t e^{\lambda(e^t - 1)} + \lambda^2 e^{2t} e^{\lambda(e^t - 1)}$$

$$E[X] = M_X'(0) = \lambda$$

$$E[X^2] = M_X''(0) = \lambda + \lambda^2$$

$$\text{VAR}(X) = E[X^2] - (E[X])^2 = \lambda$$



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

Example 2

Let $X \sim \text{Bernoulli}(p)$ distribution with probability mass
function $p(x) = \begin{cases} 1-p, & \text{if } x=0 \\ p, & \text{if } x=1 \end{cases}$.

$$\text{Then } M_X(t) = E[e^{tx}] = e^{tx} \Big|_{x=0} (1-p) + e^{tx} \Big|_{x=1} p = 1-p + e^t p.$$

Example 3

Let $X \sim \text{Binomial}(p)$ distribution.

$$M_X(t) = (1-p + e^t p)^n$$



Center for Research and Training in Innovative
Techniques of Applied Mathematics in Engineering



Department of Applied Mathematics,
Faculty of Applied Sciences
University Politehnica of Bucharest



universidad
SANJORGE



ThinkBS: Basic Sciences in Engineering Education, Erasmus Plus Project,

INNOVATIVE MATHEMATICAL MODELING TECHNIQUES: FRACTIONAL CALCULUS, WAVELET ANALYSIS, AND ESTIMATING OF NONLINEARITIES

Compound distribution

Let be X_1, X_2, \dots, X_m random variables i.i.d. that describe the same distribution function X and m becomes an integer positive and N random variable

$$\text{For } S = \sum_{i=1}^N X_i, \text{ one find: } M(S) = M(X) M(N) \\ D^2(S) = M^2(X) \cdot D^2(N) + D^2(X) M(N)$$

If $N \sim Po(\lambda) \Rightarrow M(N) = D^2(N) = \lambda; M(N^2) = \lambda^2 + \lambda$, then:

$$M(S) = \lambda M(X), D^2(S) = \lambda M(X^2).$$