


# Smooth Functions on a Manifold

## Definition (Smooth functions)

Let  $M$  be a manifold of dimension  $n$ .

- A function  $f : M \rightarrow \mathbb{R}$  is said to be  $C^\infty$  or *smooth at a point*  $p \in M$  when there is a chart  $(U, \phi)$  about  $p$  in  $M$  such that the function  $f \circ \phi^{-1} : \phi(U) \rightarrow \mathbb{R}$  is  $C^\infty$  at  $\phi(p)$  (here  $\phi(U)$  is an open subset of  $\mathbb{R}^n$ ).
- We say that  $f$  is  $C^\infty$  on  $M$  when it is  $C^\infty$  at every point of  $M$ .



2 / 23

# Smooth Functions on a Manifold

## Remark

- The smoothness condition is independent of the choice of the chart  $(U, \phi)$ .
- If  $(V, \psi)$  is another chart about  $p$  and  $f \circ \phi^{-1}$  is  $C^\infty$ , then  $f \circ \psi^{-1} = (f \circ \phi^{-1}) \circ (\phi \circ \psi^{-1})$  is  $C^\infty$  at  $p$  as well, since the transition map  $\phi \circ \psi^{-1}$  is a  $C^\infty$ .

## Remark

- If a function  $f : M \rightarrow \mathbb{R}$  is  $C^\infty$  at  $p$ , then it is automatically continuous at  $p$ .
- If  $(U, \phi)$  is a chart about  $p$  and  $f \circ \phi^{-1}$  is  $C^\infty$  at  $\phi(p)$ , then  $f = (f \circ \phi^{-1}) \circ \phi$  is continuous at  $p$ , since  $\phi$  is a continuous map.
- Therefore, any  $C^\infty$ -function on  $M$  is continuous.

3 / 23

# Smooth Functions on a Manifold

## Proposition (Proposition 6.3)

Let  $f : M \rightarrow \mathbb{R}$  be a function. Then TFAE:

- 1  $f$  is  $C^\infty$  on  $M$ .
- 2 For every chart  $(U, \phi)$  on  $M$ , the function  $f \circ \phi^{-1} : \phi(U) \rightarrow \mathbb{R}$  is  $C^\infty$ .

4 / 23


# Smooth Maps Between Manifolds

In what follows  $M$  is a manifold of dimension  $m$  and  $N$  is a manifold of dimension  $n$ .

## Definition (Smooth maps between manifolds)

Let  $F : N \rightarrow M$  be a continuous map.

- We say that  $F$  is  $C^\infty$  or *smooth* at  $p \in N$  when there are a chart  $(U, \phi)$  about  $p$  in  $M$  and a chart  $(V, \psi)$  about  $F(p)$  on  $N$  such that the map  $\psi \circ F \circ \phi^{-1} : \phi(F^{-1}(V) \cap U) \rightarrow \mathbb{R}^m$  is  $C^\infty$  at  $\phi(p)$  (here  $\phi(F^{-1}(V) \cap U)$  is an open set in  $\mathbb{R}^n$ ).
- Then map  $F$  is  $C^\infty$  on  $N$  when it is  $C^\infty$  at every point  $p \in N$ .



5 / 23

# Smooth Maps Between Manifolds

## Remark

- We assume  $F : N \rightarrow M$  to be continuous to ensure that  $F^{-1}(V)$  is an open set in  $N$ .
- When  $M = \mathbb{R}^m$  the continuity assumption can be dropped.

## Proposition (Remark 6.6)

A map  $F : N \rightarrow N$  is  $C^\infty$  at  $p$  if and only if there is a chart  $(U, \phi)$  about  $p$  in  $N$  such that the map  $F \circ \phi^{-1} : \phi(U) \rightarrow \mathbb{R}^m$  is  $C^\infty$  at  $p$  (here  $\phi(U)$  is an open set in  $\mathbb{R}^n$ ).

6 / 23

# Smooth Maps Between Manifolds

## Proposition (Proposition 6.7)

Suppose that  $F : N \rightarrow M$  is  $C^\infty$  at  $p$ . Then, for every chart  $(U, \phi)$  about  $p$  in  $N$  and every chart  $(V, \psi)$  about  $F(p)$  in  $M$ , the map  $\psi \circ F \circ \phi^{-1} : \phi(F^{-1}(V) \cap U) \rightarrow \mathbb{R}^m$  is  $C^\infty$  at  $\phi(p)$ .

## Proposition (Proposition 6.8)

Let  $F : N \rightarrow M$  be a continuous map. TFAE:

- 1  $F$  is a  $C^\infty$  map.
- 2 For every chart  $(U, \phi)$  on  $N$  and every chart  $(V, \psi)$  on  $M$ , the map  $\psi \circ F \circ \phi^{-1} : \phi(F^{-1}(V) \cap U) \rightarrow \mathbb{R}^m$  is  $C^\infty$ .

7 / 23

## Proposition (Proposition 6.9; Composition of $C^\infty$ maps)

If  $F : N \rightarrow M$  and  $G : P \rightarrow N$  are  $C^\infty$  maps (where  $P$  is a manifold), then the composition  $F \circ G : P \rightarrow M$  is a  $C^\infty$  map.

8 / 23

## Diffeomorphisms

### Definition

We say that a map  $F : N \rightarrow M$  is a *diffeomorphism* when it is a bijective  $C^\infty$  map with  $C^\infty$  inverse  $F^{-1}$ .

### Proposition (Proposition 6.10)

If  $(U, \phi)$  is a chart on  $M$ , then the coordinate map  $\phi : U \rightarrow \phi(U) \subset \mathbb{R}^m$  is a diffeomorphism.

### Proposition (Proposition 6.11)

Let  $U$  be an open subset of  $M$ . If  $F : U \rightarrow F(U) \subset \mathbb{R}^n$  is a diffeomorphism onto an open subset of  $\mathbb{R}^n$ , then the pair  $(U, F)$  is a chart on  $M$ .

9 / 23

## Smoothness in Terms of Components

### Proposition (Propositions 6.12 & 6.13)

Let  $F : N \rightarrow \mathbb{R}^m$  be a map with components  $F^1, \dots, F^m : N \rightarrow \mathbb{R}$  (so that  $F(p) = (F^1(p), \dots, F^m(p))$ ). Then TFAE:

- 1  $F$  is a  $C^\infty$ -map.
- 2 For every chart  $(U, \phi)$  on  $N$ , the map  $F \circ \phi^{-1} : \phi(U) \rightarrow \mathbb{R}^m$  is  $C^\infty$ .
- 3 All the components  $F^1, \dots, F^m : N \rightarrow \mathbb{R}$  are  $C^\infty$  maps

### Remark

We don't need to assume  $F$  to be continuous, since the 2nd and 3rd properties both imply that  $F$  is continuous.

10 / 23

## Smoothness in Terms of Components

### Proposition (Propositions 6.15 & 6.16)

Let  $F : N \rightarrow M$  be a continuous map. Then TFAE:

- 1  $F$  is a  $C^\infty$  map.
- 2 For every chart  $(V, \psi)$  on  $M$  the vector-valued function  $\psi \circ F : F^{-1}(V) \rightarrow \mathbb{R}^m$  is  $C^\infty$ .
- 3 For every chart  $(V, \psi) = (V, y^1, \dots, y^m)$  the component functions  $y^i \circ F : F^{-1}(V) \rightarrow \mathbb{R}$  are  $C^\infty$ .

### Remark

We assume  $F$  to be continuous to insure that in the 2nd and 3rd properties  $F^{-1}(V)$  is an open subset of  $N$ .

11 / 23

## Examples of Smooth Maps

### Example (Example 6.17 + Exercise 6.18)

Let  $M_1$  and  $M_2$  be manifolds.

- 1 The 1st factor projection  $\pi_1 : M_1 \times M_2 \rightarrow M_1$ ,  $\pi_1(p_1, p_2) = p_1$  is a  $C^\infty$  map. Likewise, the 2nd factor projection  $\pi_2 : M_1 \times M_2 \rightarrow M_2$  is a smooth map.
- 2 Given a manifold  $N$ , a map  $f : N \rightarrow M_1 \times M_2$  is  $C^\infty$  if and only if the components  $\pi_i \circ f : N \rightarrow M_i$  are  $C^\infty$  maps.

12 / 23

## Examples of Smooth Maps

### Example (Example 6.19)

Let  $\mathbb{S}^n \subset \mathbb{R}^{n+1}$  be the unit sphere. If  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  is a  $C^\infty$  function, then the restriction  $f|_{\mathbb{S}^n} : \mathbb{S}^n \rightarrow \mathbb{R}$  is a  $C^\infty$  function on  $\mathbb{S}^n$ .

13 / 23

## Examples of Smooth Maps

### Definition (Lie Groups)

A *Lie group* is a group  $G$  equipped with a differentiable structure such that:

- (i) The multiplication map  $\mu : G \times G \rightarrow G, (x, y) \rightarrow xy$  is a  $C^\infty$  map.
- (ii) The inverse map  $\iota : G \rightarrow G, x \rightarrow x^{-1}$  is a  $C^\infty$  map.

### Examples

- 1 The Euclidean spaces  $\mathbb{R}^n$  and  $\mathbb{C}^n$  are Lie groups under addition.
- 2 The set of non-zero complex numbers  $\mathbb{C}^\times := \mathbb{C} \setminus 0$  is a Lie group under multiplication.
- 3 The unit circle  $\mathbb{S}^1 \subset \mathbb{C}^\times$  is a Lie group under multiplication.
- 4 If  $G_1$  and  $G_2$  are Lie groups, then their Cartesian product  $G_1 \times G_2$  is again a Lie group.

14 / 23

## Examples of Smooth Maps

### Example (Example 6.21; see Tu's book)

We saw in Section 5 that the general groups  $GL(n, \mathbb{R})$  and  $GL(n, \mathbb{C})$  are manifolds. They are also Lie groups under multiplication of matrices.

### Remark

Further examples of Lie groups are studied in Section 15.

15 / 23

# Partial Derivatives

In what follows  $M$  is a manifold of dimension  $n$ .

## Reminder

If  $(U, \phi) = (U, x^1, \dots, x^n)$  a chart on  $M$ , then by definition the components  $x^1, \dots, x^n$  of  $\phi$  are given by  $x^i = r^i \circ \phi : U \rightarrow \mathbb{R}$ .

## Definition

Let  $f : M \rightarrow \mathbb{R}$  be a  $C^\infty$  function. For  $p \in U$  the *partial derivative of  $f$  with respect to  $x^i$  at  $p$*  is

$$\frac{\partial f}{\partial x^i}(p) := \frac{\partial(f \circ \phi^{-1})}{\partial r^i}(\phi(p)).$$

## Remark

The partial derivative  $\frac{\partial f}{\partial x^i}(p)$  is also denoted  $\frac{\partial}{\partial x^i} \Big|_p f$ .

16 / 23

# Partial Derivatives

## Remark

As  $\phi^{-1}(\phi(p))$  the equality  $\frac{\partial f}{\partial x^i}(p) = \frac{\partial(f \circ \phi^{-1})}{\partial r^i}(\phi(p))$  can be rewritten as

$$\frac{\partial f}{\partial x^i} \circ \phi^{-1}(\phi(p)) = \frac{\partial(f \circ \phi^{-1})}{\partial r^i}(\phi(p)).$$

Thus, as functions on  $\phi(U)$  we have

$$\frac{\partial f}{\partial x^i} \circ \phi^{-1} = \frac{\partial(f \circ \phi^{-1})}{\partial r^i}.$$

In particular, this shows that  $\frac{\partial f}{\partial x^i} : U \rightarrow \mathbb{R}$  is  $C^\infty$  function on  $U$ .

## Proposition (Proposition 6.22)

If  $(U, x^1, \dots, x^n)$  is a chart on  $M$ , then  $\frac{\partial x^i}{\partial x^j} = \delta_j^i$ .

17 / 23



In what follows  $M$  is a manifold of dimension  $m$  and  $N$  is a manifold of dimension  $n$ .

## Definition (Jacobian matrices and Jacobian determinants)

Let  $F : M \rightarrow N$  be a  $C^\infty$  map. Let  $(U, \phi) = (U, x^1, \dots, x^n)$  be a chart on  $N$  and  $(V, \psi) = (V, y^1, \dots, y^n)$  a chart on  $M$  such that  $F(U) \subset V$ . Denote  $F^i := y^i \circ F = r^i \circ \psi \circ F : U \rightarrow \mathbb{R}$  the  $i$ -th component of  $F$  in the chart  $(V, \psi)$ .

- 1 The matrix  $[\partial F^i / \partial x^j]$  is called the *Jacobian matrix* of  $F$  relative to the charts  $(U, \phi)$  and  $(V, \psi)$ .
- 2 When  $m = n$  the determinant  $\det [\partial F^i / \partial x^j]$  is called the *Jacobian determinant* of  $F$  relative to the charts.

## Remark

The Jacobian determinant is also denoted  $\partial(F^1, \dots, F^n) / \partial(x^1, \dots, x^n)$ .

18 / 23

## Remark

If  $N = U$  is an open subset of  $\mathbb{R}^n$  and  $M = V$  is an open subset of  $\mathbb{R}^m$ , and we use the charts  $(U, r^1, \dots, r^n)$  and  $(V, r^1, \dots, r^m)$ , then the Jacobian matrix  $[\partial F^i / \partial r^j]$  is the usual Jacobian matrix from calculus.

## Example (Example 6.24; Jacobian matrix of a transition map)

Let  $(U, \phi) = (U, x^1, \dots, x^n)$  and  $(V, \psi) = (V, y^1, \dots, y^n)$  be overlapping charts on  $N$ . The transition map  $\psi \circ \phi^{-1} : \phi(U \cap V) \rightarrow \psi(U \cap V)$  is a diffeomorphism between open subsets of  $\mathbb{R}^n$ . Given any  $p \in U \cap V$ , we have

$$\frac{\partial y^i}{\partial x^j}(p) = \frac{(\psi \circ \phi^{-1})^i}{\partial r^j}(\phi(p)).$$

# The Inverse Function Theorem

In what follows  $M$  and  $N$  are manifolds of dimension  $n$ .

## Reminder

By Proposition 6.11, given an open  $U \subset M$ , any diffeomorphism  $F : U \subset F(U) \subset \mathbb{R}^n$  defines a coordinate system on  $U$ , i.e.,  $(U, F)$  is a chart on  $M$ .

## Definition

We say that a  $C^\infty$  map  $F : N \rightarrow M$  is *locally invertible* or is *local diffeomorphism* near  $p \in N$  if there is an open neighborhood  $U$  of  $p$  in  $N$  such that  $F|_U : U \rightarrow F(U)$  is a diffeomorphism.

## Remark

If  $F = (F^1, \dots, F^n) : N \rightarrow \mathbb{R}^n$  is locally invertible near  $p \in N$ , then it defines a coordinate system about  $p$ .

20 / 23

# The Inverse Function Theorem

Theorem (Theorem 6.25, Inverse Function Theorem for  $\mathbb{R}^n$ ; see also Appendix B)

Let  $F = (F^1, \dots, F^n) : W \rightarrow \mathbb{R}^n$  be a  $C^\infty$ -map, where  $W$  is an open set in  $\mathbb{R}^n$ . Given any  $p \in W$ , TFAE:

- (i)  $F$  is locally invertible near  $p$ .
- (ii) The Jacobian determinant  $\det[\partial F^i / \partial x^j(p)]$  is non-zero.

21 / 23

# The Inverse Function Theorem

## Theorem (Theorem 6.26, Inverse Function Theorem for manifolds)

Let  $F : N \rightarrow M$  be a  $C^\infty$ -map. Given any  $p \in N$ , TFAE:

- (i)  $F$  is locally invertible near  $p$ .
- (ii) We have a non-zero Jacobian determinant  $\det[\partial F^i / \partial x^j(p)]$ .

## Remarks

- 1 In (ii) the Jacobian determinant  $\det[\partial F^i / \partial x^j(p)]$  relatively to some chart  $(U, x^1, \dots, x^n)$  about  $p$  in  $N$  and some chart  $(V, y^1, \dots, y^n)$  about  $F(p)$  in  $M$  and we have  $F^i = y^i \circ F$ .
- 2 The condition  $\det[\partial F^i / \partial x^j(p)] \neq 0$  is independent of the choice of the charts.

22 / 23

# The Inverse Function Theorem




Fig. 6.4. The map  $F$  is locally invertible at  $p$  because  $\psi \circ F \circ \phi^{-1}$  is locally invertible at  $\phi(p)$ .

## Corollary (Corollary 6.27)

Let  $F = (F^1, \dots, F^n) : U \rightarrow \mathbb{R}^n$  be  $C^\infty$  map on a neighborhood  $U$  of a point  $p$  in  $N$ . TFAE:

- 1  $F = (F^1, \dots, F^n)$  defines a coordinate system near  $p$ .
- 2  $\det[\partial F^i / \partial x^j(p)] \neq 0$ .

23 / 23